

Date palm seeds

- Description
- Nutritional aspects
- Nutritional tables
- References

Click on the "Nutritional aspects" tab for recommendations for ruminants, pigs, poultry, rabbits, horses, fish and crustaceans



Common names

Date seeds, dates pits, date kernels, date stones, date pips

Species

Phoenix dactylifera L. [Arecaceae]

Feed categories

- Other plant by-products
- Plant products and by-products

Related feed(s)

- Date molasses
- Date palm leaves and date pedicels
- Date palm fruits

Description

Date seeds are the by-product of date stoning, either for the production of pitted dates or for the manufacture of date paste (Ecocrop, 2011; Barreveld, 1993). The date seed is a hard coated seed, usually oblong, ventrally grooved, with a small embryo. Date pits weigh 0.5 g to 4 g and represent 6 to 20% of the fruit weight depending on maturity, variety and grade (Ecocrop, 2011; Daghir, 2008; Zaid et al., 2002; Barreveld, 1993; Göhl, 1982).

Date seeds are traditionally used for animal feed. They can also be used as a source of oil (which has antioxidant properties valuable in cosmetics), as a coffee substitute, as a raw material for activated carbon or as an adsorbent for dye-containing waters (Lecheb, 2010; Banat et al., 2003; Barreveld, 1993). Date by-products are usually fed to animals during winter, though they can be used at any time of the year (Genin et al., 2004).

Distribution

Date by-products are available in countries of production (see **Distribution** in the [Date palm fruits](#) datasheet). Date seeds are available near to where dates are packed or processed. The date industry is the main source of date pits, but they are also produced at farm level in date paste production units (Barreveld, 1993).

Processes

Date pits have a hard seed coat that makes the seed components difficult to digest. It is necessary to process the seeds before feeding them to livestock.

Soaking

Traditionally, farmers soak date pits in water for 72 hours to increase their feeding value. However this method does not improve animal intake (Barreveld, 1993).

Grinding

Grinding increases nutrient availability by breaking and removing the seed coat. However, it is energy-consuming and causes wear and tear to the machinery (Barreveld, 1993). It is recommended to first crush the seed with a disk crusher and then grind them with a hammer-mill (Barreveld, 1993; Göhl, 1982).

Chemical treatments


Alkali treatments increase the digestibility of fibrous materials. Application of a 9.6% NaOH solution to ground date pits decreased NDF content and increased *in vitro* digestion rates. This treatment was more effective on finely ground date pits (4 mm vs. 8 mm) (Al-Yousef et al., 1986).

Biological treatments

Germinated date seeds (after 74 days) have a similar chemical composition to dry pits but are softer and can be more easily eaten by animals (Sumianah et al., 1984 cited by Barreveld, 1993).

Environmental impact

The by-products of date production have always been recycled by farmers and local populations, a practice that has been considered as "an eloquent example of integrated sustainable use of renewable material resources" (El-Mously, 2001). There



is a long tradition of using cull dates and date pits to feed animals, and the crop residues serve as raw materials to create household items, furniture and building materials ([El-Mously, 2001](#)).

Datasheet citation

Heuzé V., Tran G., Delagarde R., Bastianelli D., Lebas F., 2015. *Date palm seeds*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/687> Last updated on May 11, 2015, 14:34

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

● Gilles Tran, AFZ ● Gilles Tran, AFZ

[+](#) Share / Save [f](#) [t](#) [r](#)



Date palm seeds

- Description
- Nutritional aspects
- Nutritional tables
- References

Nutritional attributes

Date seeds are a low protein feed, with about 5-11% DM crude protein. Oil content is in the 4-14% DM range. Date pits contain high and variable quantities of fibre, with a high level of lignification: crude fibre 16-51% DM, NDF 58-90% DM, ADF 41-46% DM and ADL 4-18% (Feedipedia, 2011; Boudechiche et al., 2009; Dagher, 2008; Lecheb, 2010; Barreveld, 1993; Genin et al., 2004). Varietal differences are significant (Boudechiche et al., 2009). Date pits contain appreciable amounts of K followed by P, Mg and Ca, but have a low Na content. Of the micro elements present, Fe, Mn, Zn and Cu are the most important (Barreveld, 1993).

Date seeds are tough and need to be processed before being fed to animals.

Potential constraints

Pesticide residues

Dates are fumigated to kill insects, often with methyl bromide (CH₃Br) (Glasner et al., 2002). The maximum residue level for methyl bromide on dried fruits was set at 2 mg/kg by the Codex Alimentarius (Codex Alimentarius, 2011). Other pesticides are used, including carbamates and organophosphates, that may leave residues (Khan et al., 2001).

In organic date palm production, carbon dioxide is used instead of methyl bromide (Glasner et al., 2002).

Steroid compounds

The presence of steroid compounds in date pits, notably estrone, progesterone and estriol, has been known since the 1950s, though the actual effects of these compounds on sheep growth and reproduction have yet to be clearly demonstrated (Barreveld, 1993; El-Gasim et al., 1995; El-Din et al., 2001). Treatments with hexane and diethyl ether (combined in some cases with germination) have been proposed in order to decrease the amount of steroids (El-Din et al., 2001).

Ruminants

The nutritive value of date seeds has been quite extensively studied due to their widespread availability in the countries where date production is important. It should be noted that most of the research concerning the utilization of date pits in ruminants is relatively old, not readily accessible and, therefore, difficult to assess. The renewed interest in these products is already resulting in new research work and this datasheet will be updated once enough contemporary literature becomes available.

Ground date seeds can be used up to 75% of the diet of ruminants provided that a good protein supplement (such as cottonseed cake) or urea is added. They are also useful to balance a diet that is too rich in protein, such as young pasture (Al-Wash et al., 1982). The crude protein and crude fat of date seeds are low but not negligible, and the seeds need to be processed so that the hard seed coat is no longer an obstacle to their digestion (See Processes above) (Barreveld, 1993).

Digestibility

The nutritive values of date seeds reported in the literature are variable, with *in vivo* DM digestibility in sheep ranging from 58% to 70% (El Shazly et al., 1963; Al-Yousef et al., 1993). Organic matter digestibility values higher than 80% have been reported (Richter et al., 1956; Al-Kinani et al., 1975 cited by Al-Wash et al., 1982). *In vitro* DM digestibility using the rumen fluids of goats, sheep and dromedaries were found to be much lower for dromedaries and sheep (30-35%) than for goats (52-60%) (Genin et al., 2004). Protein digestibility was generally low (less than 40%) or not measurable (Al-Yousef et al., 1993; Al-Wash et al., 1982; El Shazly et al., 1963). Date pits have a higher value than leaves and pedicels (Al-Yousef et al., 1993).

Cattle

In 2011, there was no recent research on the use of date pits in cattle feeding, even though such utilization is attested to in the literature for dairy and beef cattle (Al-Wash et al., 1982). The daily gain of calves was not affected when the percentage of date pits in the diet increased from 30 to 60% in a concentrate mixture of wheat bran, barley and sesame meal (Farhan et al., 1969 cited by Al-Wash et al., 1982).

Sheep

The optimal level of date pits in sheep diets is disputed, probably due to the large variety of experimental protocols used by researchers. In a fattening trial where lambs were given 0 to 75% date pits, the largest gain was obtained at the 75% level (with 25% alfalfa hay) (Al-Kinani et al., 1975 cited by Al-Wash et al., 1982). Another trial found that daily gain and carcass traits were improved when 50% crushed date pits were supplemented with urea and incorporated in a mixture containing alfalfa, a concentrate and molasses (Younis et al., 1981 cited by Al-Wash et al., 1982). An inclusion rate of 30% date pits gave the largest weight gain in sheep fed *Atriplex halimus* hay and a concentrate. The isonitrogenous replacement of barley grain by ground date pits, included at up 45% in the diet had no effect on total feed intake and *in vivo* DM digestibility despite a high increase in the NDF and ADF content of the diet. Fibre digestibility increased with the inclusion of date pits, which suggests that fibre digestibility of date pits was higher than that of barley grain. Feeding date pits may reduce the occurrence of acidosis more effectively than barley (Al-Owaimer et al., 2011).

Dromedaries

Raw date pits are an excellent slow release energy feed for camels during long desert journeys (Barreveld, 1993).

Pigs

Poultry

<http://www.feedipedia.org/node/687>[08/12/2016 17:37:39]

Date palm seeds

- Description
- Nutritional aspects
- Nutritional tables
- References

Tables of chemical composition and nutritional value

- Date palm (*Phoenix dactylifera*), pits

Avg: average or predicted value; SD: standard deviation; Min: minimum value; Max: maximum value; Nb: number of values (samples) used

Date palm (*Phoenix dactylifera*), pits



Main analysis	Unit	Avg	SD	Min	Max	Nb
Dry matter	% as fed	90.5	0.9	89.3	92.3	17
Crude protein	% DM	6.4	0.6	5.8	8.4	21
Crude fibre	% DM	28.9	10.1	15.7	47.9	19
NDF	% DM	66.1	4.5	57.5	69.4	6
ADF	% DM	45.5	3.9	41.2	52.8	6
Lignin	% DM	8.5	6.0	3.9	18.4	6
Ether extract	% DM	6.7	1.4	4.9	9.0	18
Ash	% DM	1.9	0.5	1.3	3.2	20
Gross energy	MJ/kg DM	20.0				*

Minerals	Unit	Avg	SD	Min	Max	Nb
Calcium	g/kg DM	3.5	2.0	0.6	6.7	14
Phosphorus	g/kg DM	2.3	0.6	1.3	3.3	14

Amino acids	Unit	Avg	SD	Min	Max	Nb
Alanine	% protein	4.2	0.1	4.1	4.4	4
Arginine	% protein	4.1	3.9	1.1	9.8	4
Aspartic acid	% protein	7.4	1.4	6.3	9.1	4
Cystine	% protein	1.7		1.7	1.8	2
Glutamic acid	% protein	13.3	3.5	9.6	17.2	4
Glycine	% protein	3.8	0.9	3.0	4.7	4
Histidine	% protein	1.5	0.6	0.9	2.1	4
Isoleucine	% protein	3.0	0.3	2.8	3.5	4
Leucine	% protein	5.7	0.1	5.6	5.8	4
Lysine	% protein	3.8	1.0	2.9	4.7	4
Methionine	% protein	1.6	0.1	1.5	1.6	4
Phenylalanine	% protein	3.5	0.1	3.4	3.6	4
Proline	% protein	3.0	0.3	2.7	3.3	4
Serine	% protein	3.8	0.3	3.5	4.0	4
Threonine	% protein	3.2	0.2	3.0	3.4	4
Tryptophan	% protein	0.8				1
Tyrosine	% protein	1.3	0.2	1.1	1.4	3
Valine	% protein	3.9	0.9	3.1	4.9	4

Ruminant nutritive values	Unit	Avg	SD	Min	Max	Nb
OM digestibility, Ruminant	%	71.4				1
Energy digestibility, ruminants	%	70.2				*
DE ruminants	MJ/kg DM	14.0				*
ME ruminants	MJ/kg DM	11.5				*
Nitrogen digestibility, ruminants	%	18.0		0.0	36.0	2

Pig nutritive values	Unit	Avg	SD	Min	Max	Nb
Energy digestibility, growing pig	%	44.8				*
DE growing pig	MJ/kg DM	8.9				*



Poultry nutritive values	Unit	Avg	SD	Min	Max	Nb
TME poultry	MJ/kg DM	3.3				1

The asterisk * indicates that the average value was obtained by an equation.

References

AFZ, 2011; Al-Yousef et al., 1993; El Shazly et al., 1963; El-Gasim et al., 1986; El-Hag et al., 1992; Fangauf et al., 1957; Genin et al., 2004; Hussein et al., 1998; Kamel et al., 1981; Kamel et al., 1981; Vandepopuliere et al., 1995

Last updated on 24/10/2012 00:44:07

Datasheet citation

Heuzé V., Tran G., Delagarde R., Bastianelli D., Lebas F., 2015. *Date palm seeds*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/687> *Last updated on May 11, 2015, 14:34*

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

- Gilles Tran, AFZ
- Gilles Tran, AFZ



Date palm seeds

Description	Nutritional aspects	Nutritional tables	References
-------------	---------------------	--------------------	------------

References

Aboul-Ela, S. ; El-Hindawy, M. ; Sherif, S. Y. ; Tawfik, E. S. ; Attia, A. I., 1999. Evaluating date pits as a waste product of food industries in feeding NZW rabbits. Cahiers Options Méditerranéennes, 41: 57-65 / In: Testik A., Baselga M. (eds.). 2nd International Conference on rabbit production in hot climates, Zaragoza, CIHEAM-IAMZ, 1998/09/07-09, Adana (Turkey)

Afifi, M. ; Abdou, F. ; El-Sayed, M., 1966. Date stone meal as a substitute for barley in chick rations. Trop. Agric. (Trinidad), 43:167

Al-Asghar, N.A., 1988. Date palm seeds as food for carp (*Cyprinus carpio* L.). J. Coll. Sci. King Saud Univ., 19 (1): 59-64

Al-Bowait, M. ; Al-Sultan, S. I., 2006. Aspects of the serum biochemistry, carcass quality and organoleptic characteristics of broilers fed alkali-treated date pits. Int. J. Poult. Sci., 5 (3): 284-288

Al-Bowait, M. ; Al-Sultan, S. I., 2007. The effect of partial replacement of maize by alkali-treated date pits on broiler growth, survival rate and economic returns. Livest. Res. Rural Dev., 19 (9): 121

Al-Kinani, L. M.; Al-Wash, A. M., 1975. Study of different proportions of date stones in the ration for fattening Awassi lambs. Iraqi J. Agric. Sci., 10: 53-62

Al-Owaimer, A. N. ; El-Waziry, A. M. ; Koohmaraie, M. ; Zahran, S. M., 2011. The use of ground date pits and *Atriplex halimus* as alternative feeds for sheep. Aust. J. Basic Appl. Sci., 5 (5): 1154-1161

Al-Wash, A. H. ; DePeters, E. J., 1982. The use of date stones for feeding and fattening ruminant animals. World Review of Animal Production, 18 (3): 29-32

Al-Yousef, Y. ; Belyea, R. L. ; Vandepopuliere, J. M., 1986. Sodium hydroxide treatment of date pits. In: Proceeding of The Second Symposium on The Date Palm. King Faisal University, Al-Hassa, Saudi Arabia. March, 1986. Vol. II

Al-Yousef, Y. M. ; Al-Mulhim, F. N. ; El-Hag, G. A. ; El-Gasim, E. A., 1993. Apparent digestibility of discarded and date pits together with other agricultural by-products. Proceeding of The Third Symposium on The Date Palm. King Faisal University, Al-Hassa, Saudi Arabia. January 1993 (Rajab,1413) Vol. II

Ali, K. T. ; Fine, N. C. ; Sarsam, N. H. ; McLeroy, G. B., 1956. Macerated dates and ground date stones as feeds for fattening sheep. Empire J. Exp. Agric., 24: 323-337

Ali, B. H. ; Bashir, A. K. ; Alhadrami, G., 1999. Reproductive hormonal status of rats treated with date pits. Food Chem., 66: 437-441

Banat, F. ; Al-Asheha, S. ; Al-Makhadmeha, L., 2003. Evaluation of the use of raw and activated date pits as potential adsorbents for dye containing waters. Process Biochemistry, 39 (2): 193-202

Barreveld, W. H., 1993. Date palm products. FAO Agricultural Services Bulletin No. 101

Belal, I. E. H. ; Al-Owafeir, M. A., 2004. Incorporating date pits *Phoenix dactylifera* and their sprouts in semi-purified diets for Nile tilapia *Oreochromis niloticus* (L.). J. World Aquacult. Soc., 35 (4): 452-459

Belal, I. E. H., 2008. Evaluating fungi-degraded date pits as a feed ingredient for Nile tilapia *Oreochromis niloticus* L.. Aquacult. Nutr., 14 (5): 445-452

Boudechiche, L. ; Arab, A. ; Tahar, A. ; Ouzrout, R., 2009. Study of chemical composition of date stones for use in animal feed. Livest. Res. Rural Dev., 21 (5)

Chapoutot, P., 1998. Étude de la dégradation *in situ* des constituants pariétaux des aliments pour ruminants. Thèse Docteur en Sciences Agronomiques, Institut National Agronomique Paris-Grignon, Paris (FRA), 1998/11/17.

Codex Alimentarius, 2011. Pesticides residues in Food and feed. Methyl Bromide. Codex Alimentarius, FAO/WHO Food Standards

Daghir, N. J., 2008. Poultry production in hot climates. Second Edition, Cabi Series, CABI

Ecocrop, 2011. Ecocrop database. FAO

El Shazly, K. ; Ibrahim, E. A. ; Kara, H. A., 1963. Nutritional value of date seeds for sheep. J. Anim. Sci., 22 (4): 894-897

El-Din, A. E. M. M. N. ; El-Hameed, A. K. E. A., 2001. Study on the preparation of date seeds for animal feeding. 2nd Inter. Conf. on Date Palms. March 25-27, 2001, Al-Ain, United Arab Emirates Univ., pp. 887-898

El-Gasim, E. A. ; Al-Yousef, Y. A. ; Humeida, A. M., 1995. Possible hormonal activity of date pits and flesh fed to meat animals. Food Chem., 52: 149-152

El-Mously, H., 2001. The industrial use of the date palm residues: an eloquent example of sustainable development. 2nd Inter. Conf. on Date Palms. March 25-27, 2001, Al-Ain, United Arab Emirates Univ., pp. 866-886

FAO, 2011. FAOSTAT. Food and Agriculture Organization of the United Nations

Farhan, S. M. A. ; Jumah, H. T., 1969. The use of date byproducts for fattening calves. Iraqi J. Agric. Sci., 4: 7

Genin, D. ; Kadri, A. ; Khorchani, T. ; Sakkal, K. ; Belgacem, F. ; Hamadi, M., 2004. Valorisation of date-palm by-products (DPBP) for livestock feeding in Southern Tunisia. I-Potentialities and traditional utilisation. In: Ben Salem, H., Nefzaoui A., Morand-Fehr, P. (eds). Nutrition and feeding strategies of sheep and goats under harsh climates (Stratégies de nutrition et d'alimentation des ovins et caprins en climats rigoureux), Zaragoza : CIHEAM-IAMZ, 2004: 227-232

Glasner, B. B. ; Botes, A. ; Zaid, A. ; Emmens, J., 2002. Date harvesting, packinghouse management and marketing aspects. In: Zaid, A. Arias-Jimenez, E. J. Date palm cultivation. FAO plant production and protection paper, 156 rev.1. FAO, Rome

Göhl, B., 1982. Les aliments du bétail sous les tropiques. FAO, Division de Production et Santé Animale, Roma, Italy

Hermes, I. H. ; Al-Homidan, A. H., 2004. Effects of using date waste (whole dates and date pits) on performance, egg components and quality characteristics of Baladi Saudi and Leghorn laying hens.. Egyptian J. Nutr. Feeds

Hussein, A. S. ; Alhadrami, G. A. ; Khalil, Y. H., 1998. The use of dates and date pits in broiler starter and finisher diets. Bioresource Technol., 66 (3): 219-223

Hussein, A.S. ; Alhadrami, G.A., 2003. Effect of Enzyme Supplementation and Diets Containing Date Pits on Growth and Feed Utilization of Broiler Chicks. J. Aric. Marine Sci., 8 (2): 67-71

Ibrahim, M. R. ; El-Banna, H. M. ; El-Manylawi, M. A., 2010. Evaluating utilization of ground date stone with or without Kemzyme in the diets of growing New Zealand rabbits. IV International Date Palm Conference, Acta Hort. (ISHS), 882 : 691-697

Kamel, B. S. ; Diab, M. F. ; Ilian, M. A. ; Salman, A. J., 1981. Nutritional value of whole dates and date pits in broiler rations. Poult. Sci., 60 (5): 1005-1011

Khan, A. J. ; Azam, K. M. ; Razvi, S. A., 2001. Pesticide residue analysis of date palm fruits by gas chromatography mass spectrophotometry. 2nd Inter. Conf. on Date Palms. March 25-27, 2001, Al-Ain, United Arab Emirates Univ., pp. 211-215

Lecheb, F., 2010. Extraction et caractérisation physico-chimique et biologique de la matière grasse du noyau des dattes: essai d'incorporation dans une crème cosmétique de soin. Mémoire de magister, Spécialité Génie alimentaire, Faculté des Sciences de l'Ingénieur, Université, M'Hamed Bougara, Boumerdès

Mahmoud, H. A., 2008. Using date seed meal in Dandarawi laying hens and growing cockerels diets.. Assiut Vet. Med. J., 54: 11-23

Masoudi, A. ; Bojarpour, M. ; Chaji, M. ; Eslami, M. ; Mirzadeh, K., 2010. Economic value of date pits replaced with maize in broiler chicken Diet. J. Anim. Vet. Adv., 9 (11): 1578-1581

Nany, S. I. ; Soheir, K. ; Zeidan, M. ; Deraz, T. A. ; Afify, A. A., 2009. Productive and reproductive performance of lactating buffaloes fed on rations containing different levels of date stone. Proceedings of The 2nd Scientific Conf. of Animal Wealth Research in the Middle East & North Africa. Cairo International Convention Center, Egypt, 24 – 26 October, (2009) pp. 415- 426

Richter, K. von; Becker, M., 1956. Composition and nutritional value of dates and date parts as a result of trials on ruminants and pigs. Zeitschrift für Tierernährung und Futtermittelkunde, 11: 289

Sumianah, Gh. M. ; Makki, Y. M. ; Rumney, T. G., 1984. Changes in the chemical composition of three cultivars of date seed during germination. Date Palm J., 3 (2)

Vandepopuliere, J. M. ; Al-Yousef, Y. ; Lyons, J. J., 1995. Dates and date pits as ingredients in broiler starting and Coturnix quail breeder diets. Poult. Sci., 74 (7): 1134-1142

Younis, R. A. ; Al-Wash, A. H., 1981. The effect of different levels of urea and date stones in the ration on its digestion and performance of Awassi lambs. Iraqi J. Agric. Sci., ?

Zaghari, M. ; Taherkhani, R. ; Ghasemi, M. ; Shivazad, M., 2009. Estimation of metabolisable energy content of date pit and its effect on lipid and protein oxidation in broiler chicks. J. Sci. Food Agric., 89 (13): 2336-2341

Zaid, A. ; Arias-Jimenez, E. J., 2002. Date palm cultivation. FAO Plant production and protection paper 156, Rev. 1. FAO, Rome

46 references found

Datasheet citation

Heuzé V., Tran G., Delagarde R., Bastianelli D., Lebas F., 2015. *Date palm seeds*. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <http://www.feedipedia.org/node/687> Last updated on May 11, 2015, 14:34

English correction by Tim Smith (Animal Science consultant) and Hélène Thiollet (AFZ)

Image credits

Gilles Tran, AFZ Gilles Tran, AFZ

Share / Save

